

A MODEL OF COMPOSITE QUARKS AND LEPTONS

T. PRADHAN

Institute of Physics, Bhubaneswar-751005, INDIA

ABSTRACT

In continuation of our model of composite electron, its companion neutrinos, photon and electro-weak gauge bosons as well as their super-partners based on extended Lorentz group $SU(3) \otimes SU(3)$ in (8+1) dimensional space-time, we show that the remaining leptons, quarks, lepto-quarks and gluons as well as their respective super-partners can be considered as composites of fundamental particles belonging to the $5 \otimes 5$ dimensional representation of the further extended Lorentz group $SU(5) \otimes SU(5)$ in (24+1) dimensional space-time. The fundamental particles constituting the composite ones do not possess quantum numbers of the latter such as charge, hypercharge etc. Since according to our estimate based on experimental limit on the electron life-time, the mass of the fundamental particles is in the Planckian regime, the composite particles would break up into their constituents at these energies and lose their quantum numbers.

In an earlier communication [1], we showed that the electron, its companion neutrino and the electro-weak gauge-bosons as well as their super-partners are composites of fundamental particles belonging to the $3 \otimes 3$ dimensional representation of the extended Lorentz group $SU(3) \otimes SU(3)$ in (8+1) dimensional space-time which are held together by forces provided by gauge-particles belonging to the $8 \otimes 8$ dimensional representation. The fine structure constant of the gauge coupling was found to be unity from the vanishing of the neutrino mass and an estimate of $m > 10^{22}$ GeV for the mass of the fundamental particles was obtained from the experimental limit $\tau > 10^{23}$ years for the life-time of the electron indicating thereby that the composite nature of electron, photon etc will be revealed at Planckian energies. In this communication we extend this model so as to encompass all leptons, quarks, lepto-quarks, gluons and their super-partners in addition to particles covered in the previous model. It is shown that this is accomplished through the extended $SU(5) \otimes SU(5)$ Lorentz group in (24+1) dimensional space-time.

The Dirac equations in spinor representation (symmetrical representation) in (24+1) dimensional space-time reads:

$$\begin{aligned} (p_0 - \frac{1}{4}\Lambda_a p_a)\xi &= m\eta \\ (p_0 + \frac{1}{4}\Lambda_a p_a)\eta &= m\xi \end{aligned} \tag{1}$$

a = 1,2,.....24 where

$$\xi = \begin{pmatrix} \xi_1 \\ \xi_2 \\ . \\ . \\ \xi_5 \end{pmatrix} \quad \eta = \begin{pmatrix} \eta_1 \\ \eta_2 \\ . \\ . \\ \eta_5 \end{pmatrix}$$

and Λ_a are $SU(5)$ matrices. This pair of equations split into the four pairs of equations (2) and (3) in (8+1) dimensional space-times:

$$\begin{aligned} (p_0 - \frac{1}{2}\lambda_i p_i)\xi^{(\alpha)} &= m\eta^{(\alpha)} \\ (p_0 + \frac{1}{2}\lambda_i p_i)\eta^{(\alpha)} &= m\xi^{(\alpha)} \end{aligned} \tag{2}$$

where

$$\xi^\alpha = \begin{pmatrix} \xi_1 \\ \xi_2 \\ \xi_\alpha \end{pmatrix} \quad \eta^\alpha = \begin{pmatrix} \eta_1 \\ \eta_2 \\ \eta_\alpha \end{pmatrix}$$

$$\alpha = 3, 4, 5$$

$$i = \begin{cases} 1, 2, 3, 4, 5, 6, 7, 8 & \text{for } \alpha = 3 \\ 1, 2, 3, 9, 10, 13, 14, 24 & \text{for } \alpha = 4 \\ 1, 2, 3, 11, 12, 15, 16, 24 & \text{for } \alpha = 5 \end{cases}$$

and

$$\begin{aligned} (p_0 - \frac{1}{2}\lambda_i p_i)\xi^{(c)} &= m\eta^{(c)} \\ (p_0 + \frac{1}{2}\lambda_i p_i)\eta^{(c)} &= m\xi^{(c)} \end{aligned} \tag{3}$$

where

$$\xi^{(c)} = \begin{pmatrix} \xi_4 \\ \xi_5 \\ \xi_3 \end{pmatrix} \quad \eta^{(c)} = \begin{pmatrix} \eta_4 \\ \eta_5 \\ \eta_3 \end{pmatrix}$$

$$i = 17, 18, 19, 20, 21, 22, 23, 24$$

Each of these pairs of equations can be further reduced to ten pairs of (3+1) space-time equations (4), (5), (6) and (7):

$$\begin{aligned} (p_0 - \vec{\sigma} \cdot \vec{p})\xi_p &= m\eta_p \\ (p_0 + \vec{\sigma} \cdot \vec{p})\eta_p &= m\xi_p \end{aligned} \tag{4}$$

where

$$\begin{aligned} \xi_p &= \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix} \quad \eta_p = \begin{pmatrix} \eta_1 \\ \eta_2 \end{pmatrix} \\ \vec{p} &= (p_1, p_2, p_3) \\ (p_0 - \vec{\sigma} \cdot \vec{\pi}^{(\alpha)})\xi_\pi^{(\alpha)} &= m\eta_\pi^{(\alpha)} \\ (p_0 + \vec{\sigma} \cdot \vec{\pi}^{(\alpha)})\eta_\pi^{(\alpha)} &= m\xi_\pi^{(\alpha)} \end{aligned} \tag{5}$$

where

$$\xi_\alpha^{(\pi)} = \begin{pmatrix} \xi_1 \\ \xi_\alpha \end{pmatrix} \quad \alpha = 3, 4, 5$$

$$\begin{aligned}
\vec{\pi}^{(3)} &= (p_4, p_5, \frac{p_8}{\sqrt{3}}), \quad \vec{\pi}^{(4)} = (p_9, p_{10}, \frac{p_{24}}{\sqrt{15}}), \quad \vec{\pi}^{(5)} = (p_{15}, p_{16}, \frac{p_{24}}{\sqrt{15}}) \\
(p_0 - \vec{\sigma} \cdot \vec{\kappa}^{(\alpha)}) \xi_{\kappa}^{(\alpha)} &= m \eta_{\kappa}^{(\alpha)} \\
(p_0 + \vec{\sigma} \cdot \vec{\kappa}^{(\alpha)}) \eta_{\kappa}^{(\alpha)} &= m \xi_{\kappa}^{(\alpha)}
\end{aligned} \tag{6}$$

where

$$\begin{aligned}
\xi_{\kappa}^{(\alpha)} &= \begin{pmatrix} \xi_2 \\ \xi_{\alpha} \end{pmatrix} \quad \alpha = 3, 4, 5 \\
\vec{\kappa}^{(3)} &= (p_6, p_7, \frac{p_8}{\sqrt{3}}), \quad \vec{\kappa}^{(4)} = (p_{13}, p_{14}, \frac{p_{24}}{\sqrt{15}}), \quad \vec{\kappa}^{(5)} = (p_{11}, p_{12}, \frac{p_{24}}{\sqrt{15}}) \\
(p_0 - \vec{\sigma} \cdot \vec{p}^{(r)}) \xi_r &= m \eta_r \\
(p_0 + \vec{\sigma} \cdot \vec{p}^{(r)}) \eta_r &= m \xi_r
\end{aligned} \tag{7}$$

where

$$\begin{aligned}
\xi_r &= \begin{pmatrix} \xi_4 \\ \xi_5 \end{pmatrix} \quad \eta_r = \begin{pmatrix} \eta_4 \\ \eta_5 \end{pmatrix} \\
\vec{p}_r &= (p_{21}, p_{22}, p_{23})
\end{aligned}$$

and identical equations for $(\xi^{(b)}, \eta^{(b)})$ and $(\xi^{(g)}, \eta^{(g)})$ with

$$\begin{aligned}
\xi_b &= \begin{pmatrix} \xi_4 \\ \xi_3 \end{pmatrix}, \quad \eta_b = \begin{pmatrix} \eta_4 \\ \eta_3 \end{pmatrix}, \quad \xi_g = \begin{pmatrix} \xi_5 \\ \xi_3 \end{pmatrix}, \quad \eta_g = \begin{pmatrix} \eta_5 \\ \eta_3 \end{pmatrix} \\
\vec{p}_b &= (p_{19}, p_{20}, \frac{p_{24}}{\sqrt{15}}), \quad \vec{p}_g = (p_{17}, p_{18}, \frac{p_{24}}{\sqrt{15}})
\end{aligned}$$

Construction of composite particles can be made with the aid of scalars and spinors obtained from the ten fundamental fields

$$\begin{aligned}
\xi_p &= \begin{pmatrix} \xi_1 \\ \xi_2 \end{pmatrix}, \quad \xi_{\pi}^{(\alpha)} = \begin{pmatrix} \xi_1 \\ \xi_{\alpha} \end{pmatrix}, \quad \xi_{\kappa}^{(\alpha)} = \begin{pmatrix} \xi_2 \\ \xi_{\alpha} \end{pmatrix} \\
\alpha &= 3, 4, 5 \\
\xi_r &= \begin{pmatrix} \xi_4 \\ \xi_5 \end{pmatrix}, \quad \xi_b = \begin{pmatrix} \xi_4 \\ \xi_3 \end{pmatrix}, \quad \xi_g = \begin{pmatrix} \xi_5 \\ \xi_3 \end{pmatrix}
\end{aligned}$$

the equations of motion of which are given in eqns(4), (5), (6) and (7). The composition of various particles is as follows:

Leptons:

$$l^{(\alpha)} = \begin{pmatrix} e & \mu & \tau \\ \nu_e & \nu_\mu & \nu_\tau \end{pmatrix} = \xi_p \begin{pmatrix} \xi_\pi^{(\alpha)+} \\ \xi_\kappa^{(\alpha)+} \end{pmatrix} \quad (8)$$

Quarks :

$$q_i^{(\alpha)} = \begin{pmatrix} u_i & c_i & t_i \\ d_i & s_i & b_i \end{pmatrix} = \Phi_{p_i} \begin{pmatrix} \xi_\pi^{(\alpha)+} \\ \xi_\kappa^{(\alpha)+} \end{pmatrix} \quad (9)$$

where $\Phi_{p_i} = \xi_p \xi_i^+$ is a first rank spinor in the (p_0, p_1, p_2, p_3) space.

Electro-weak Gauge Bosons :

$$A = \Psi_p \phi_\pi \quad Z = \Psi_p \phi_\kappa \quad W = \Psi_p \phi_{\pi\kappa} \quad (10)$$

where $\Psi_p = \xi_p \xi_p^+$ is a second rank spinor and

$$\phi_\pi = \frac{1}{\sqrt{3}} \sum_\alpha \xi_\pi^{(\alpha)+} \xi_\pi^{(\alpha)}, \quad \phi_\kappa = \frac{1}{\sqrt{3}} \sum_\alpha \xi_\kappa^{(\alpha)+} \xi_\kappa^{(\alpha)}, \quad \phi_{\pi\kappa} = \frac{1}{\sqrt{3}} \sum_\alpha \xi_\pi^{(\alpha)+} \xi_\kappa^{(\alpha)}$$

are scalars.

Gluons :

$$g_{ij} = \Psi_p \phi_{ij}, \quad i, j = r, b, g \quad (11)$$

where $\phi_{ij} = \xi_i^+ \xi_j$ is a scalar.

Lepto-quarks :

$$X_i = \Psi_p \phi_\pi \xi_i, \quad Y_i = \Psi_p \phi_\kappa \xi_i \quad (12)$$

In all these constructions we have only the ξ components of the particles concerned. Their η components can be obtained by parity reversal. Further, their spin states will be obtained from those of ξ_p .

The super-partners of all the above particles can be constructed as follows:

Scalar Leptons :

$$\tilde{l}^{(\alpha)} = \begin{pmatrix} \tilde{e}, & \tilde{\mu} & \tilde{\tau} \\ \tilde{\nu}_e & \tilde{\nu}_\mu & \tilde{\nu}_\tau \end{pmatrix} = \phi_p \begin{pmatrix} \xi_\pi^{(\alpha)+} \\ \xi_\kappa^{(\alpha)+} \end{pmatrix} \quad (13)$$

where $\phi_p = \xi_p^+ \xi_p$ is a scalar.

Quarks :

$$\tilde{q}_i^{(\alpha)} = \begin{pmatrix} \tilde{u}_i & \tilde{c}_i & \tilde{t}_i \\ \tilde{d}_i & \tilde{s}_i & \tilde{b}_i \end{pmatrix} = \phi_i \begin{pmatrix} \xi_\pi^{(\alpha)+} \\ \xi_\kappa^{(\alpha)+} \end{pmatrix} \quad (14)$$

where $\phi_i = \phi_p \xi_i$ is a scalar.

Gluinos :

$$\tilde{g}_{ij} = \xi_p \phi_{ij} \quad (15)$$

Gauginos:

$$\tilde{A} = \xi_p \phi_\pi \quad \tilde{Z} = \xi_p \phi_\kappa \quad \tilde{W} = \xi_p \phi_{\pi\kappa} \quad (16)$$

As stated earlier, the gauge particles that provide the glue belong to the regular representation of $SU(5) \otimes SU(5)$. The dynamics of these particles will be similar to those of the $SU(3) \otimes SU(3)$ case discussed in our previous communication, but more complex. We shall take up this problem in a later publication.

We conclude by remarking that the fundamental particles of our model do not possess quantum numbers like charge, hypercharge etc which are attributes of the composite particles. The latter will break up into the constituent fundamental particles at Planckian energies and lose their quantum numbers.

References

1. T. Pradhan, Institute of Physics Preprint IP/BBSR/97-46.